

Chouinard, Luc

- Kramer, S. M.*, May, J. R.*, Patrick, D. J.*, **Chouinard, L.**, Boyer, M., Doyle, N., et al. (2009). Instilled or injected purified natural capsaicin has no adverse effects on rat hindlimb sensory-motor behavior or osteotomy repair. *Anesthesia and Analgesia*, 109(1), 249-257.
- Rosset, P.*, & **Chouinard, L. E.** (2009). Characterization of site effects in montreal, canada. *Natural Hazards*, 48(2), 295-308.
- Oestergaard, S.*, Rasmussen, K. E.*, Doyle, N., Varela, A.*, **Chouinard, L.**, Smith, S. Y., et al. (2008). Evaluation of cartilage and bone degradation in a murine collagen antibody-induced arthritis model. *Scandinavian Journal of Immunology*, 67(3), 304-312.
- Oestergaard, S.*, **Chouinard, L.**, Doyle, N., Smith, S. Y., Tankó, L. B.*, & Qvist, P.* (2006). Early elevation in circulating levels of C-telopeptides of type II collagen predicts structural damage in articular cartilage in the rodent model of collagen-induced arthritis. *Arthritis and Rheumatism*, 54(9), 2886-2890.
- Oestergaard, S.*, **Chouinard, L.**, Doyle, N., Karsdal, M. A.*, Smith, S. Y., Qvist, P.*, et al. (2006). The utility of measuring C-terminal telopeptides of collagen type II (CTX-II) in serum and synovial fluid samples for estimation of articular cartilage status in experimental models of destructive joint diseases. *Osteoarthritis and Cartilage*, 14(7), 670-679.